

CONSENT FOR COLONOSCOPY

Purpose of Examination	Colonoscopy is the examination to diagnose colon polyps, colon cancers, inflammatory diseases in your bowl by using a colonoscope. It also takes biopsies and removes polyps.
Cautions	1) An anticoagulant or antiplatelet drug such as Aspirin, Ticlopedine, Warfarin can cause to increase the risk of bleeding when taking biopsies or removing polyps. If you take these medications, you should consult your doctor and stop them one week before the examination day. 2) Do not take insulin shots in the morning of examination after consulting your doctor. 3) Do not take a tablet for diabetes in the morning of the examination. 4) If you take an antihypertensive drug or an anticonvulsant, take it with a minimum amount of water in the morning of the examination.
Possible Complications	You might have some abdominal pain and bloating after the colonoscopy. Rarely, perforation to the bowl (0.03%), peritonitis, bleeding can happen. If you feel that your abdomen becomes tense, or experience symptoms including severe abdominal pain, continuous bloody stool, diarrhea, dizziness, nausea, vomiting, pallor and cold sweating you must contact us immediately or go to the nearest emergency room because you might need an emergency treatment or surgery.

Cases which require cautions during endoscopy (Answer correctly)

	Yes	No
Fast	<input type="checkbox"/>	<input type="checkbox"/>
Hypersensitiveness to medications or food (Antibiotic, anesthetic, egg, bean, sulfites, sulfonamide)	<input type="checkbox"/>	<input type="checkbox"/>
Cardiovascular diseases (Angina pectoris, myocardial infection, arrhythmia)	<input type="checkbox"/>	<input type="checkbox"/>
Kidney diseases (Chronic glomerulonephritis, renal failure)	<input type="checkbox"/>	<input type="checkbox"/>
Liver diseases (Chronic hepatitis, alcoholic hepatitis, hepatic cirrhosis)	<input type="checkbox"/>	<input type="checkbox"/>
Respiratory diseases (Sleep apnea, asthma, chronic obstructive pulmonary disease, severe snoring, a cold)	<input type="checkbox"/>	<input type="checkbox"/>
Benign prostatic hyperplasia	<input type="checkbox"/>	<input type="checkbox"/>
Glaucoma	<input type="checkbox"/>	<input type="checkbox"/>
Hypertension	<input type="checkbox"/>	<input type="checkbox"/>
Thyroid diseases	<input type="checkbox"/>	<input type="checkbox"/>
Anticoagulant medications (Aspirin, Coumadin, Warfarin)	<input type="checkbox"/>	<input type="checkbox"/>
Pregnancy / Possibility to pregnancy / Breast feeding	<input type="checkbox"/>	<input type="checkbox"/>

Colonoscopy History	
Surgery History (e.g. abdominal surgery)	

During the examination, a biopsy and polypectomy might be needed for an accurate diagnosis. It will incur additional costs.

I have been fully informed about the necessity, nature, possible complications of the colonoscopy, and I fully understand risks of a complication beyond human control or any unexpected accidents caused by idiosyncrasy. I hereby consent to performing the colonoscopy.

	Date:	(mm/dd/yyyy)
	Examinee:	(Signature)
Relationship to the examinee:	Parent/Guardian:	(Signature)



Guide to take PLENVU®

My colonoscopy

Date :

Time :

Name :


How to reconstitute PLENVU®



Dose1 [mango flavor]

- 1 Pour the contents of Dose1 sachet into 1000mL bottle.
- 2 Add 1000mL of **cold-water**.
- 3 Shake bottle until powder is completely dissolved.

※ PLENVU® can be refrigerated, but must be taken within 24 hours after it is mixed with water.



Dose2 [fruit punch flavor]

- 1 Pour the contents of Dose2 both sachet A and sachet B into 1000mL bottle.
- 2 Add 1000mL of **cold-water**.
- 3 Shake bottle until powder is completely dissolved.

How to drink PLENVU®

- 1 Sip reconstituted PLENVU very slowly within next 1 hour.

Dose1



Take reconstituted PLENVU®

drink 500mL of water

Dose2



Take reconstituted PLENVU®

drink 500mL of water

Take the
PharmaSimethicone

※ You can drink water or tea before, during and after taking PLENVU®. But you must stop taking any fluids 3 hour before your procedure.

For Morning Colonoscopy

Take Dose 1

Date: The evening before colonoscopy

Time : PM 9:00

Take Dose 2

Date : The next morning

Time :

Drink at least 4 hours before your
visiting hospital

For Afternoon Colonoscopy

Take Dose 1

Date : On the day of your colonoscopy

Time : AM 6:00

Take Dose 2

Date : On the day of your colonoscopy

Time : AM 9:00



Food Intake

Precautions

- Please be consulted if you are treated because of heart failure or kidney disease or you are treated with pill or insulin because of diabetes.
- Let your doctor know If you are taking anticoagulants(asprin, warfarin, coumadin, ticlopidin or clopidogrel, etc.). Do not take it within a week before you take PLENVU®.
- In case polypectomy is performed, going abroad can be possible only after at least one week after polypectomy.
- If you have constipation you need dietary control before one week and you might need to take laxative.
- It's important to take PLENVU® completely to clean your bowel before the procedure.
- If you take reexamine, you should pay an extra charge

3 days prior to procedure

Do not eat fruits w/seeds, sesames or mushrooms.

Do not eat various beans, vegetables, seaweeds or Kimchi.

Do eat Egg, Tofu, Fish, etc.



1 day prior to procedure



For Morning Colonoscopy : you can grab a light breakfast and lunch.
(polished rice soup+soy sauce polished rice+soup without solid, sponge cake)

For Afternoon Colonoscopy : you can grab a light lunch and dinner.
(polished rice soup+soy sauce polished rice+soup without solid, sponge cake)



한국의료재단 | IFC
Korea Medical Foundation | 종합검진센터